

Properties of the Bethe Ansatz equations for Richardson-Gaudin models

Inna Lukyanenko, Phillip Isaac, Jon Links

*Centre for Mathematical Physics,
School of Mathematics and Physics,
The University of Queensland*




THE UNIVERSITY
OF QUEENSLAND
AUSTRALIA

Some History

- ▶ **Quantum Inverse Scattering Method (QISM)** for twisted periodic boundary conditions [Faddeev, Kulish, Sklyanin, Takhtajan 1979].
- ▶ **BCS model** of superconductivity [Bardeen, Cooper and Schrieffer 1957].
 - ▶ Solved for the **rational case** [Richardson 1963].
 - ▶ **Integrals of motion** for the Richardson model [Cambiaggio et al. 1997].
 - ▶ **Eigenvalues** for the Richardson model [Sierra 2000].
 - ▶ Generalized to the **trigonometric case** using Gaudin's method [Amico et al. 2001], [Dukelsky et al. 2001].
 - ▶ Reformulated through the **quasi-classical limit** of QISM [Zhou et al. 2002], [von Delft and Poghossian 2002].
- ▶ **Some extensions:** [Ovchinnikov 2003], [Dunning and Links 2004], [Ibañez et al. 2009], [Skrypnik 2009], [Dukelsky et al. 2010, 2011], [Links and Marquette 2013].

Some History


- ▶ **Boundary QISM (BQISM)** for open-boundary conditions, for the case of XXZ spin chain [Sklyanin 1988].



- ▶ What is the effect of the “boundary” for **Richardson-Gaudin models**?
 - ▶ **Gaudin magnet** with boundary [Hikami 1995].
 - ▶ **Quasi-classical limit** of the BQISM [Di Lorenzo et al. 2002].
 - ▶ **Generalized** Gaudin systems [Skrypnik 2006, 2007, 2010].
 - ▶ **Trigonometric** Gaudin model [Cirilio António et al. 2013].

Outline

In the quasi-classical limit (QC)



R-matrix

R-matrix is an operator $R(u) \in \text{End}(V \otimes V)$ ($V \cong \mathbb{C}^2$, $u \in \mathbb{C}$) satisfying the **Yang-Baxter equation (YBE)** in $\text{End}(V \otimes V \otimes V)$:

$$R_{12}(u - v)R_{13}(u)R_{23}(v) = R_{23}(v)R_{13}(u)R_{12}(u - v)$$

- **Rational solution** ($\eta \in \mathbb{C}$, $P(u \otimes v) = v \otimes u$, $\forall u, v \in V$)

$$R^{\text{rat}}(u) = \frac{1}{u + \eta}(uI \otimes I + \eta P) = \frac{1}{u + \eta} \begin{pmatrix} u + \eta & 0 & 0 & 0 \\ 0 & u & \eta & 0 \\ 0 & \eta & u & 0 \\ 0 & 0 & 0 & u + \eta \end{pmatrix}.$$

- **Trigonometric solution**

$$R^{\text{trig}}(u) = \frac{1}{\sinh(u + \eta)} \begin{pmatrix} \sinh(u + \eta) & 0 & 0 & 0 \\ 0 & \sinh u & \sinh \eta & 0 \\ 0 & \sinh \eta & \sinh u & 0 \\ 0 & 0 & 0 & \sinh(u + \eta) \end{pmatrix}.$$

Rational limit $\lim_{\nu \rightarrow 0} \frac{\sinh(\nu x)}{\nu} = x$:

Trigonometric \longrightarrow **Rational**

QISM [Faddeev et al. 1979]

Monodromy matrix $\in \text{End}(V_a \otimes V^{\otimes \mathcal{L}})$ (where $V_a = \mathbb{C}^2$ is the auxiliary space, $V^{\otimes \mathcal{L}} = \underbrace{V \otimes V \otimes \dots \otimes V}_{\mathcal{L} \text{ times}}$ is the quantum space, $\mathcal{L} \in \mathbb{N}$)

$$T_a(u) := \begin{pmatrix} e^{-\eta\gamma} & 0 \\ 0 & e^{\eta\gamma} \end{pmatrix} R_{a\mathcal{L}}(u - \varepsilon_{\mathcal{L}}) \dots R_{a2}(u - \varepsilon_2) R_{a1}(u - \varepsilon_1) = \begin{pmatrix} A(u) & B(u) \\ C(u) & D(u) \end{pmatrix}$$

Transfer matrix $t(u) := \text{tr}_a(T_a(u)) = A(u) + D(u) \in \text{End}(V^{\otimes \mathcal{L}})$:

$$\boxed{[t(u), t(v)] = 0 \quad \forall u, v \in \mathbb{C}}$$

Then $t(u)$ generates a set of **mutually commuting** operators $\{C_j\}$:

$$t(u) = \sum_{j=-\infty}^{\infty} C_j u^j.$$

Take any function of $\{C_j\}$ as the Hamiltonian. Then $\{C_j\}$ are mutually commuting **integrals of motion**.

Algebraic Bethe Ansatz

Start with a **reference state** $\Omega \in V^{\otimes \mathcal{L}}$:

$$B(u)\Omega = 0, \quad A(u)\Omega = a(u)\Omega, \quad D(u)\Omega = d(u)\Omega, \quad C(u)\Omega \neq 0.$$

Then (for the **trigonometric** R -matrix)

$$\Phi(v_1, \dots, v_N) = C(v_1) \dots C(v_N)\Omega$$

is an **eigenstate** of $t(u)$ with the **eigenvalue**

$$\Lambda(u, v_1, \dots, v_N) = a(u) \prod_{k=1}^N \frac{\sinh(u - v_k + \eta)}{\sinh(u - v_k)} + d(u) \prod_{k=1}^N \frac{\sinh(u - v_k - \eta)}{\sinh(u - v_k)},$$

if $\Phi \neq 0$ and v 's satisfy the **Bethe Ansatz equations (BAE)**

$$\frac{a(v_k)}{d(v_k)} = \prod_{i \neq k}^N \frac{\sinh(v_k - v_i - \eta)}{\sinh(v_k - v_i + \eta)}, \quad k = 1, \dots, N$$

For the **rational** R -matrix: $\sinh(x) \rightarrow x$.

Algebraic Bethe Ansatz

Start with a **reference state** $\Omega \in V^{\otimes \mathcal{L}}$:

$$B(u)\Omega = 0, \quad A(u)\Omega = a(u)\Omega, \quad D(u)\Omega = d(u)\Omega, \quad C(u)\Omega \neq 0.$$

Then (for the **trigonometric** R -matrix)

$$\Phi(v_1, \dots, v_N) = C(v_1) \dots C(v_N) \Omega$$

is an **eigenstate** of $t(u)$ with the **eigenvalue**

$$\Lambda(u, v_1, \dots, v_N) = a(u) \prod_{k=1}^N \frac{\sinh(u - v_k + \eta)}{\sinh(u - v_k)} + d(u) \prod_{k=1}^N \frac{\sinh(u - v_k - \eta)}{\sinh(u - v_k)},$$

if $\Phi \neq 0$ and v 's satisfy the **Bethe Ansatz equations (BAE)**

$$\frac{a(v_k)}{d(v_k)} = \prod_{i \neq k}^N \frac{\sinh(v_k - v_i - \eta)}{\sinh(v_k - v_i + \eta)}, \quad k = 1, \dots, N$$

For the **rational** R -matrix: $\sinh(x) \rightarrow x$.

Quasi-classical limit (QC)

Derive the expressions for $a(u)$ and $d(u)$ for $\Omega = \begin{pmatrix} 0 \\ 1 \end{pmatrix}^{\otimes \mathcal{L}}$:

$$a(u) = e^{-\eta\gamma} \prod_{l=1}^{\mathcal{L}} \frac{\sinh(u - \varepsilon_l - \eta/2)}{\sinh(u - \varepsilon_l)}, \quad d(u) = e^{\eta\gamma} \prod_{l=1}^{\mathcal{L}} \frac{\sinh(u - \varepsilon_l + \eta/2)}{\sinh(u - \varepsilon_l)}.$$

Then the **BAE**:

$$e^{-2\eta\gamma} \prod_{l=1}^{\mathcal{L}} \frac{\sinh(v_k - \varepsilon_l - \eta/2)}{\sinh(v_k - \varepsilon_l + \eta/2)} = \prod_{i \neq k}^N \frac{\sinh(v_k - v_i - \eta)}{\sinh(v_k - v_i + \eta)}.$$

Take first non-zero term as $\eta \rightarrow 0$:

QISM	$\xrightarrow{\eta \rightarrow 0}$	QC QISM
------	------------------------------------	---------

Quasi-classical limit (QC)

Derive the expressions for $a(u)$ and $d(u)$ for $\Omega = \begin{pmatrix} 0 \\ 1 \end{pmatrix}^{\otimes \mathcal{L}}$:

$$a(u) = e^{-\eta\gamma} \prod_{l=1}^{\mathcal{L}} \frac{\sinh(u - \varepsilon_l - \eta/2)}{\sinh(u - \varepsilon_l)}, \quad d(u) = e^{\eta\gamma} \prod_{l=1}^{\mathcal{L}} \frac{\sinh(u - \varepsilon_l + \eta/2)}{\sinh(u - \varepsilon_l)}.$$

Then the **BAE**:

$$e^{-2\eta\gamma} \prod_{l=1}^{\mathcal{L}} \frac{\sinh(v_k - \varepsilon_l - \eta/2)}{\sinh(v_k - \varepsilon_l + \eta/2)} = \prod_{i \neq k}^N \frac{\sinh(v_k - v_i - \eta)}{\sinh(v_k - v_i + \eta)}.$$

Take first non-zero term as $\eta \rightarrow 0$:

$$\boxed{\text{QISM} \xrightarrow{\eta \rightarrow 0} \text{QC QISM}}$$

- QC trig. QISM [Amico et al. 2001], [Dukelsky et al. 2001]

$$2\gamma + \sum_{l=1}^{\mathcal{L}} \coth(v_k - \varepsilon_l) = 2 \sum_{i \neq k}^N \coth(v_k - v_i) \quad (\diamond)$$


- QC rat. QISM [Richardson 1963]

$$2\gamma + \sum_{l=1}^{\mathcal{L}} \frac{1}{v_k - \varepsilon_l} = \sum_{i \neq k}^N \frac{2}{v_k - v_i} \quad (\heartsuit)$$

QC trig. QISM (\diamond) $\xrightarrow{\text{rational limit}}$ QC rat. QISM (\heartsuit)

- Change of variables $v_i = \ln y_i$, $\varepsilon_l = \ln z_l$ in (\diamond) gives

$$(N - \mathcal{L}/2 + \gamma - 1) + \sum_{l=1}^{\mathcal{L}} \frac{y_k^2}{y_k^2 - z_l^2} = \sum_{i \neq k}^N \frac{2y_k^2}{y_k^2 - y_i^2} \quad (\diamond')$$



Reflection equations [Cherednik 1984]

Start with the **trigonometric** R -matrix. In addition to the YBE we require that it satisfies the **reflection equations** for some $K^\pm \in \text{End}(V)$, referred to as the **reflection matrices**:

$$\begin{cases} R_{12}(u-v)K_1^-(u)R_{21}(u+v)K_2^-(v) = K_2^-(v)R_{12}(u+v)K_1^-(u)R_{21}(u-v), \\ R_{12}(v-u)K_1^+(u)R_{21}(u+v)K_2^+(v) = K_2^+(v)R_{12}(u+v)K_1^+(u)R_{21}(v-u), \end{cases}$$

where $\mathcal{R}(u) \equiv R(-u - 2\eta)$. Easy to check that

$$K^-(u) = \begin{pmatrix} \sinh(\xi^- + u) & 0 \\ 0 & \sinh(\xi^- - u) \end{pmatrix},$$

$$K^+(u) = \begin{pmatrix} \sinh(\xi^+ + u + \eta) & 0 \\ 0 & \sinh(\xi^+ - u - \eta) \end{pmatrix}$$

satisfy these equations for any $\xi^\pm \in \mathbb{C}$.

BQISM [Sklyanin 1988]

Define the **double row monodromy matrix** $\in \text{End}(V_a \otimes V^{\otimes \mathcal{L}})$:

$$T_a(u) := R_{a\mathcal{L}}(u - \varepsilon_{\mathcal{L}}) \dots R_{a1}(u - \varepsilon_1) K_a^-(u) \times \\ \times R_{a1}^{-1}(-u - \varepsilon_1) \dots R_{a\mathcal{L}}^{-1}(-u - \varepsilon_{\mathcal{L}}) = \begin{pmatrix} A(u) & B(u) \\ C(u) & D(u) \end{pmatrix}.$$

The **transfer matrix**

$$t(u) := \text{tr}_a(K_a^+(u) T_a(u))$$

satisfies

$$\boxed{[t(u), t(v)] = 0 \quad \forall u, v \in \mathbb{C}}$$

Thus, it is a generating function for the **integrals of motion!**

Algebraic Bethe Ansatz

Introduce $\tilde{a}(u) = (2u)a(u) - \eta d(u)$. Start with a reference state Ω and look for other eigenstates in the form $\Phi(v_1, \dots, v_N) = C(v_1) \dots C(v_N) \Omega$.

Eigenvalues:

$$\Lambda(u, v_1, \dots, v_N) = \tilde{a}(u) \frac{\sinh(\xi^+ + u + \eta/2)}{\sinh 2u} \prod_{k=1}^N \frac{\sinh(u - v_k + \eta) \sinh(u + v_k + \eta)}{\sinh(u - v_k) \sinh(u + v_k)} +$$

$$+ d(u) \frac{\sinh(2u + \eta) \sinh(\xi^+ - u + \eta/2)}{\sinh 2u} \prod_{k=1}^N \frac{\sinh(u - v_k - \eta) \sinh(u + v_k - \eta)}{\sinh(u - v_k) \sinh(u + v_k)},$$

BAE:

$$\frac{\tilde{a}(v_k)}{d(v_k) \sinh(2v_k - \eta)} \frac{\sinh(\xi^+ + v_k + \eta/2)}{\sinh(\xi^+ - v_k + \eta/2)} = \prod_{i \neq k}^N \frac{\sinh(v_k - v_i - \eta) \sinh(v_k + v_i - \eta)}{\sinh(v_k - v_i + \eta) \sinh(v_k + v_i + \eta)}$$

Quasi-classical limit (QC)

If we substitute $\eta = 0$ the BAE will take the following form:

$$\frac{\sinh(\xi^- + v_k) \sinh(\xi^+ + v_k)}{\sinh(\xi^- - v_k) \sinh(\xi^+ - v_k)} = 1.$$

Choose $\xi^- = \xi^-(\eta)$, $\xi^+ = \xi^+(\eta)$, so that this holds as $\eta \rightarrow 0$.

Take

$$\xi^+ = \eta\alpha, \quad \xi^- = \eta\beta$$

• QC trig. BQISM

$$\begin{aligned} & -2(\alpha + \beta + 1) \coth v_k + \sum_{l=1}^{\mathcal{L}} (\coth(v_k - \varepsilon_l) + \coth(v_k + \varepsilon_l)) = \\ & = 2 \sum_{i \neq k}^N (\coth(v_k - v_i) + \coth(v_k + v_i)) \end{aligned}$$



- **QC rat. BQISM**

$$-(\alpha + \beta + 1) + \sum_{l=1}^{\mathcal{L}} \frac{v_k^2}{v_k^2 - \varepsilon_l^2} = \sum_{i \neq k}^N \frac{2v_k^2}{v_k^2 - v_i^2} \quad (\spadesuit)$$


Same expression as (\diamond') **QC trig. QISM** with the change of variables!

$$\text{QC rat. BQISM } (\spadesuit) \Leftrightarrow \text{QC trig. QISM } (\diamond)$$

- **QC trig. BQISM**

Change of variables $v_k = \ln y_k$, $\varepsilon_l = \ln z_l$ in (\clubsuit) gives

$$\begin{aligned} &-(\alpha + \beta + 1) \frac{y_k^2 + 1}{y_k^2 - 1} + \sum_{l=1}^{\mathcal{L}} \left(\frac{y_k^2}{y_k^2 - z_l^2} + \frac{1}{y_k^2 z_l^2 - 1} \right) = \\ &= \sum_{i \neq k}^N \left(\frac{2y_k^2}{y_k^2 - y_i^2} + \frac{2}{y_k^2 y_i^2 - 1} \right) \end{aligned} \quad (\clubsuit')$$



Introduce an **additional parameter** ρ : $v_k \rightarrow v_k + \frac{\rho}{2}$, $\varepsilon_l \rightarrow \varepsilon_l + \frac{\rho}{2}$.

QC rat. BQISM (♠) $\xrightarrow{\rho \rightarrow \infty}$ **QC rat. QISM (♡)**

$$-(\alpha + \beta + 1) + \sum_{l=1}^{\mathcal{L}} \frac{v_k^2 + \rho v_k + \rho^2/4}{v_k^2 - \varepsilon_l^2 + \rho(v_k - \varepsilon_l)} = 2 \sum_{i \neq k}^N \frac{v_k^2 + \rho v_k + \rho^2/4}{v_k^2 - v_i^2 + \rho(v_k - v_i)}.$$

Rescale $\alpha + \beta = \rho(\alpha' + \beta')/4$ and consider $\rho \rightarrow \infty$:

$$-(\alpha' + \beta') + \sum_{l=1}^{\mathcal{L}} \frac{1}{v_k - \varepsilon_l} = \sum_{i \neq k}^N \frac{2}{v_k - v_i}.$$


QC trig. BQISM (♣') $\xrightarrow{\rho \rightarrow \infty}$ **QC trig. QISM (◇')** Set $\rho/2 = \ln \sigma$:

$$-(\alpha + \beta + 1) \frac{\sigma^2 y_k^2 + 1}{\sigma^2 y_k^2 - 1} + \sum_{l=1}^{\mathcal{L}} \left(\frac{y_k^2}{y_k^2 - z_l^2} + \frac{1}{\sigma^4 y_k^2 z_l^2 - 1} \right) = \sum_{i \neq k}^N \left(\frac{2y_k^2}{y_k^2 - y_i^2} + \frac{2}{\sigma^4 y_k^2 y_i^2 - 1} \right)$$


and consider $\sigma \rightarrow \infty$:

$$-(\alpha + \beta + 1) + \sum_{l=1}^{\mathcal{L}} \frac{y_k^2}{y_k^2 - z_l^2} = \sum_{i \neq k}^N \frac{2y_k^2}{y_k^2 - y_i^2}.$$

Thus, we have



But in fact



Variable change

Substitute $v_k \rightarrow y_k - y_k^{-1}$, $\varepsilon_l \rightarrow z_l - z_l^{-1}$ into ():

$$-(\alpha + \beta + 1) \frac{y_k + y_k^{-1}}{y_k - y_k^{-1}} + \sum_{l=1}^{\mathcal{L}} \frac{y_k^2 - y_k^{-2}}{y_k^2 + y_k^{-2} - z_l^2 - z_l^{-2}} = 2 \sum_{i \neq k}^N \frac{y_k^2 - y_k^{-2}}{y_k^2 + y_k^{-2} - y_i^2 - y_i^{-2}}.$$

Using

$$\frac{y_k^2 - y_k^{-2}}{y_k^2 + y_k^{-2} - z_l^2 - z_l^{-2}} = \frac{y_k^2}{y_k^2 - z_l^2} + \frac{1}{y_k^2 z_l^2 - 1}$$

we obtain ():

$$-(\alpha + \beta + 1) \frac{y_k^2 + 1}{y_k^2 - 1} + \sum_{l=1}^{\mathcal{L}} \left(\frac{y_k^2}{y_k^2 - z_l^2} + \frac{1}{y_k^2 z_l^2 - 1} \right) = \sum_{i \neq k}^N \left(\frac{2y_k^2}{y_k^2 - y_i^2} + \frac{2}{y_k^2 y_i^2 - 1} \right).$$

QC rat. BQISM () $\xrightarrow{\text{variable change}}$ **QC trig. BQISM ()**

Thank you for your attention!

